

OFFICE INFORMATION:

PROCEDURE DATE:

ARRIVAL TIME:

LOCATION:

Preparing for Your Colonoscopy With OsmoPrep®

OsmoPrep is a medication used to clean your colon before colonoscopy. Colonoscopy is the #1 prevention tool for colorectal cancer.¹ The success of your colonoscopy depends on how you prepare for the procedure. Be sure to take the following steps:

1

Follow your physician's instructions regarding medicines to avoid and diet to follow before your procedure.

2

Take the entire OsmoPrep bowel prep regimen as described on the following page.

- It is important to take *all* of the medication and liquids so that your doctor will be able to see your entire colon clearly
- Be sure to stay hydrated. Drink extra clear liquids before and after your colonoscopy

3

Make arrangements to have someone drive you home after the procedure.

- Colonoscopy involves sedation, and you will not be allowed to leave unaccompanied

OsmoPrep is available at most retail pharmacy outlets. If your pharmacy does not have OsmoPrep in stock, it can be ordered and delivered within 24 hours.

Please see www.TabletPrep.com for additional information including a medication guide for OsmoPrep.


OsmoPrep
(sodium phosphate monobasic monohydrate, USP
and sodium phosphate dibasic anhydrous, USP) Tablets
The Tablet Prep

IMPORTANT SAFETY INFORMATION

There have been rare, but serious reports of acute phosphate nephropathy in patients who received oral sodium phosphate products for colon cleansing prior to colonoscopy. Some cases have resulted in permanent impairment of renal function and some patients required long-term dialysis. While some cases have occurred in patients without identifiable risk factors, patients at increased risk of acute phosphate nephropathy may include those with increased age, hypovolemia, increased bowel transit time (such as bowel obstruction), active colitis, or baseline kidney disease, and those using medicines that affect renal perfusion or function (such as diuretics, angiotensin converting enzyme [ACE] inhibitors, angiotensin receptor blockers [ARBs], and possibly nonsteroidal anti-inflammatory drugs [NSAIDs]).

It is important to use the dose and dosing regimen as recommended (PM/AM split dose).

Please see full Prescribing Information for OsmoPrep, including **BOXED WARNINGS**.

OsmoPrep® (sodium phosphate monobasic monohydrate, USP, and sodium phosphate dibasic anhydrous, USP) Tablets are indicated for cleansing of the colon as a preparation for colonoscopy in adults 18 years of age or older.

Considerable caution should be advised before OsmoPrep is used in patients with severe renal insufficiency, congestive heart failure, ascites, unstable angina, gastric retention, ileus, severe chronic constipation, bowel perforation, toxic megacolon, gastric bypass or stapling surgery, or hypomotility syndrome. Use with caution in patients with impaired renal function, patients with a history of seizures or at higher risk of seizure, patients with higher risk of cardiac arrhythmias, known or suspected electrolyte disturbances (such as dehydration), or people taking drugs that affect electrolyte levels. Patients with electrolyte abnormalities such as hypernatremia, hyperphosphatemia, hypokalemia, or hypocalcemia should have their electrolytes corrected before treatment with OsmoPrep.

OsmoPrep is contraindicated in patients with a known allergy or hypersensitivity to sodium phosphate salts or any of its ingredients, and in patients with biopsy-proven acute phosphate nephropathy. In clinical trials, the most commonly reported adverse reactions (reporting frequency >3%) were abdominal bloating, nausea, abdominal pain, and vomiting. It is recommended that patients receiving OsmoPrep be advised to adequately hydrate before, during, and after the use of OsmoPrep.


OsmoPrep® Dosing Instructions

OsmoPrep dosing consists of 2 series of tablets taken with a clear liquid of your choice.

Beginning at 3pm-day before procedure

First Dosing Series


One dose (swallow 4 tablets with 8 oz of any clear liquid) every 15 minutes for a total of 5 doses (20 tablets)



Beginning at 6 hours before check-in time

Second Dosing Series

One dose (swallow 4 tablets with 8 oz of any clear liquid) every 15 minutes for a total of 3 doses (12 tablets)



Examples of clear liquids are water, ginger ale, apple juice, sports drink, and lemonade (no pulp). Do not drink any liquids colored red or purple. If you are not sure, ask your doctor if a particular drink is acceptable.

You must complete the *entire* OsmoPrep regimen to help ensure the most effective cleansing. This will be 32 tablets. If you do not complete the regimen, your doctor may not be able to see your colon clearly.

You may have clear liquids only on the day of the exam, up to 4 hours before your check-in time.

Importance of Hydration

A colonoscopy prep will cause the body to lose a lot of fluid. This can result in illness due to the loss of too much body fluid (dehydration). It's important that you prepare your body by drinking extra clear liquids before the prep. Stay hydrated by drinking clear liquids during the prep as directed. Replenish your system by drinking clear liquids after returning home from your colonoscopy.

Please see front side for Important Safety Information.
Call your doctor for medical advice if you experience side effects.

If you have any questions, please call our office at _____

Reference: 1. Rex DK, Johnson DA, Anderson JC, Schoenfeld PS, Burke CA, Iradimi JM. American College of Gastroenterology guidelines for colorectal cleansing. 2008. *Am J Gastroenterol*. 2009;104:739-750.



Web site: www.salix.com 1700 Perimeter Park Drive, Morrisville, NC 27560 Tel • 866.669.SLXP (7597)
©2009 Salix Pharmaceuticals, Inc. All rights reserved. Printed in USA. OSM 09/14

